

東京大学大学院工学系研究科 総合研究機構  
第18回「次世代電子顕微鏡法」講演会

2024年2月2日 15:00 – 16:00 ハイブリッド開催

## Beyond vibrational spectroscopy: hunting the signature of elusive quasiparticles with monochromated STEM-EELS



Demie Kepatsoglou<sup>1,2</sup>, Quentin M. Ramasse<sup>1,3</sup>

<sup>1</sup>SuperSTEM Laboratory, Daresbury

<sup>2</sup>Department of Physics, Engineering and Technology, University of York, UK

<sup>3</sup>Schools of Physics, Astronomy & Chemical and Process Engineering, University of Leeds, UK

Engineering the structural or chemical architecture of functional materials at the nano or even atomic level enables emergent properties that rely on the interplay between fundamental properties of matter such as charge, local atomic-scale chemistry, and recently spin. Methodological developments in high-resolution energy-loss spectroscopy have made it possible to study these phenomena at near-atomic scale in the scanning transmission electron microscope. For instance, recent results exploiting the energy loss near-edge structure, which arises from core-level excitations, pave the way towards mapping electronic orbitals, such as  $\pi^*$  and  $\sigma^*$  states in epitaxial graphene, in real space. [1] In the low energy loss regime, a dark field EELS detection geometry is used to reveal atomic-scale variations in acoustic and optical phonon excitations in materials, while single-atom impurities can even be shown to have a characteristic vibrational response [2]. In a system consisting of  $\text{Bi}_2\text{Se}_3$  films grown by chemical vapor deposition on epitaxial graphene, observations using these techniques highlight the interplay between the various phonon modes and the Dirac plasmons in the topological insulator  $\text{Bi}_2\text{Se}_3$ , in addition to a further direct interrogation of the chemical bonds.

Finally, the prospects for observing the excitation of spin waves, or magnons, arising from the collective excitation of the electrons' spin in a lattice and which qualitatively occupy the same energy range as phonons, are explored through preliminary experiments and the development of a theoretical framework based on the diffuse scattering of electrons due to magnons. [3,4]

[1] M. Bugnet *et al.*, *Phys. Rev. Lett.* **128** (2022), 116401

[2] F. S. Hage *et al.*, *Science* **367** (2020), aba1136

[3] K. Lyon *et al.*, *Phys. Rev. B* **104** (2021), 214418

[4] J.A. Castellanos-Reyes *et al.*, *Phys. Rev. B* **108** (2023), 134435



東京大学  
次世代電子顕微鏡法  
社会連携講座

主催: 「次世代電子顕微鏡法」社会連携講座  
e-mail: [ishikawa@sigma.t.u-tokyo.ac.jp](mailto:ishikawa@sigma.t.u-tokyo.ac.jp)